

| Prevalence >=5 µg/dL by Town for 2018 in Rhode Island | | | |
|--|-------------------------------------|------------------------|-----------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source : RI CLPPP and LESS Database 2018 data as of 2/12/2019 | | | |
| TownName | # Children with BLU>=5mc g/dL | # children Screened | Prevalence Rate |
| Barrington | 1 | 463 | 0.2% |
| Bristol | 6 | 342 | 1.8% |
| Burrillville | 3 | 244 | 1.2% |
| Central Falls | 31 | 751 | 4.1% |
| Charlestown | 0 | 75 | 0.0% |
| Coventry | 6 | 585 | 1.0% |
| Cranston | 35 | 1,628 | 2.1% |
| Cumberland | 6 | 701 | 0.9% |
| East Greenwich | 2 | 257 | 0.8% |
| East Providence | 23 | 1,344 | 2.0% |
| Exeter | 0 | 66 | 0.0% |
| Foster | 1 | 55 | 1.8% |
| Glocester | 2 | 105 | 1.9% |
| Hopkinton | 5 | 125 | 4.0% |
| Jamestown | 0 | 45 | 0.0% |
| Johnston | 2 | 466 | 0.4% |
| Lincoln | 5 | 400 | 1.3% |
| Little Compton | 1 | 31 | 3.2% |
| Middletown | 3 | 303 | 1.0% |
| Narragansett | 1 | 76 | 1.3% |
| New Shoreham | 0 | 22 | 0.0% |
| Newport | 24 | 418 | 5.7% |
| North Kingstown | 5 | 430 | 1.2% |
| North Providence | 11 | 626 | 1.8% |
| North Smithfield | 4 | 175 | 2.3% |
| Pawtucket | 58 | 1,920 | 3.0% |
| Portsmouth | 0 | 274 | 0.0% |
| Providence | 316 | 6,361 | 5.0% |
| Richmond | 1 | 82 | 1.2% |
| Scituate | 3 | 169 | 1.8% |
| Smithfield | 1 | 292 | 0.3% |
| South Kingstown | 1 | 336 | 0.3% |
| Tiverton | 5 | 337 | 1.5% |
| Warren | 6 | 214 | 2.8% |
| Warwick | 13 | 1,393 | 0.9% |
| West Greenwich | 0 | 84 | 0.0% |
| West Warwick | 8 | 604 | 1.3% |
| Westerly | 3 | 283 | 1.0% |
| Woonsocket | 44 | 1,147 | 3.8% |
| Total | 636 | 23,034 | 2.8% |

| Prevalence >=6 µg/dL by Town for 2017 in Rhode Island | | | |
|--|--------------------------------|---------------------|-----------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source : RI CLPPP and LESS Database as of 1/29/2018 | | | |
| TownName | # Children with BLLs>=Smc g/dL | # children Screened | Prevalence Rate |
| Barrington | 9 | 452 | 2.0% |
| Bristol | 13 | 367 | 3.5% |
| Burrillville | 8 | 299 | 2.7% |
| Central Falls | 46 | 826 | 5.6% |
| Charlestown | 2 | 73 | 2.7% |
| Coventry | 11 | 563 | 2.0% |
| Granston | 62 | 1765 | 3.5% |
| Cumberland | 12 | 699 | 1.7% |
| East Greenwich | 4 | 289 | 1.4% |
| East Providence | 40 | 1076 | 3.7% |
| Exeter | 1 | 97 | 1.0% |
| Foster | 1 | 68 | 1.5% |
| Glocester | 3 | 143 | 2.1% |
| Hopkinton | 6 | 122 | 4.9% |
| Jamestown | 0 | 44 | 0.0% |
| Johnston | 11 | 566 | 1.9% |
| Lincoln | 13 | 429 | 3.0% |
| Little Compton | 1 | 84 | 2.9% |
| Middletown | 3 | 326 | 0.9% |
| Narragansett | 5 | 98 | 5.1% |
| New Shoreham | 1 | 23 | 4.3% |
| Newport | 25 | 497 | 5.0% |
| North Kingstown | 10 | 493 | 2.0% |
| North Providence | 14 | 658 | 2.1% |
| North Smithfield | 2 | 185 | 1.1% |
| Pawtucket | 88 | 2063 | 4.3% |
| Portsmouth | 4 | 263 | 1.5% |
| Providence | 431 | 6584 | 6.4% |
| Richmond | 2 | 74 | 2.7% |
| Scituate | 6 | 208 | 2.9% |
| Smithfield | 4 | 298 | 1.3% |
| South Kingstown | 10 | 392 | 2.6% |
| Tiverton | 11 | 332 | 3.3% |
| Unknown | 0 | 1 | 0.0% |
| Warren | 10 | 234 | 4.3% |
| Warwick | 29 | 1517 | 1.9% |
| West Greenwich | 0 | 71 | 0.0% |
| West Warwick | 13 | 619 | 2.1% |
| Westerly | 6 | 336 | 1.8% |
| Woonsocket | 36 | 1218 | 3.0% |
| Totals | 953 | 24,501 | 3.9% |

| Prevalence >=5 µg/dL by Town for 2016 in Rhode Island | | | |
|---|--------------------------------|---------------------|-----------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source: RI CLPPP and LESS Database as of 1/30/2016 | | | |
| TownName | # Children with BLI >=5mc g/dL | # children Screened | Prevalence Rate |
| Barrington | 9 | 469 | 1.9% |
| Bristol | 13 | 386 | 3.4% |
| Burrillville | 7 | 313 | 2.2% |
| Central Falls | 59 | 810 | 6.5% |
| Charlestown | 1 | 61 | 1.6% |
| Coventry | 9 | 562 | 1.6% |
| Cranston | 71 | 1,680 | 4.2% |
| Cumberland | 17 | 693 | 2.5% |
| East Greenwich | 4 | 294 | 1.4% |
| East Providence | 62 | 1,213 | 5.1% |
| Exeter | 1 | 84 | 1.2% |
| Foster | 1 | 76 | 1.3% |
| Gloucester | 3 | 143 | 2.1% |
| Hopkinton | 4 | 116 | 3.4% |
| Jamestown | 1 | 45 | 2.2% |
| Johnston | 15 | 545 | 2.8% |
| Lincoln | 9 | 411 | 2.2% |
| Little Compton | 3 | 44 | 6.8% |
| Middletown | 11 | 421 | 2.6% |
| Narragansett | 3 | 105 | 2.9% |
| New Shoreham | 4 | 20 | 20.0% |
| Newport | 24 | 526 | 4.6% |
| North Kingstown | 8 | 473 | 1.7% |
| North Providence | 20 | 715 | 2.8% |
| North Smithfield | 3 | 194 | 1.5% |
| Pawtucket | 123 | 2,029 | 6.0% |
| Portsmouth | 8 | 297 | 2.7% |
| Providence | 528 | 6,661 | 7.9% |
| Richmond | 5 | 76 | 6.6% |
| Schuyler | 4 | 175 | 2.3% |
| Smithfield | 1 | 290 | 0.3% |
| South Kingstown | 14 | 373 | 3.8% |
| Tiverton | 9 | 344 | 2.6% |
| Unknown | 0 | 4 | 0.0% |
| Warren | 17 | 278 | 6.1% |
| Warwick | 46 | 1,497 | 3.1% |
| West Greenwich | 0 | 87 | 0.0% |
| West Warwick | 27 | 640 | 4.2% |
| Westerly | 11 | 374 | 2.9% |
| Woonsocket | 45 | 1,214 | 3.7% |
| Totals | 1,192 | 24,738 | 4.8% |

Prevalence >=5 µg/dL by Town for 2015 in Rhode Island

For Children <= 72 months of Age at the time of test

Data Source :RI CLPPP and LESS Database as of 2/15/2016

| TownName | # Children with BfC>=5mc g/dL | # children Screened | Prevalence Rate |
|------------------|-------------------------------------|------------------------|-----------------|
| Barrington | 6 | 452 | 1.3% |
| Bristol | 12 | 451 | 2.7% |
| Burnsville | 16 | 313 | 5.1% |
| Central Falls | 66 | 875 | 7.5% |
| Charlestown | 4 | 72 | 5.6% |
| Coventry | 20 | 580 | 3.4% |
| Cranston | 75 | 1,683 | 4.5% |
| Cumberland | 15 | 739 | 2.0% |
| East Greenwich | 6 | 300 | 2.0% |
| East Providence | 63 | 1,251 | 5.0% |
| Exeter | 2 | 70 | 2.9% |
| Foster | 6 | 72 | 8.3% |
| Glocester | 4 | 124 | 3.2% |
| Hopkinton | 7 | 128 | 5.5% |
| Jamestown | 1 | 51 | 2.0% |
| Johnston | 4 | 543 | 0.7% |
| Lincoln | 10 | 424 | 2.4% |
| Little Compton | 3 | 46 | 6.5% |
| Middletown | 11 | 391 | 2.8% |
| Narragansett | 2 | 96 | 2.1% |
| New Shoreham | 4 | 22 | 18.2% |
| Newport | 33 | 564 | 5.9% |
| North Kingstown | 7 | 417 | 1.7% |
| North Providence | 13 | 628 | 2.1% |
| North Smithfield | 4 | 209 | 1.9% |
| Out Of State | 1 | 1 | 100.0% |
| Pawtucket | 156 | 2,299 | 6.8% |
| Portsmouth | 4 | 303 | 1.3% |
| Providence | 593 | 5,950 | 8.5% |
| Richmond | 3 | 74 | 4.1% |
| Saunderstown | 3 | 169 | 1.8% |
| Smithfield | 3 | 292 | 1.0% |
| South Kingstown | 16 | 342 | 4.7% |
| Tiverton | 8 | 343 | 2.3% |
| Unknown | 0 | 4 | 0.0% |
| Warren | 12 | 268 | 4.5% |
| Warwick | 37 | 1,471 | 2.5% |
| West Greenwich | 0 | 85 | 0.0% |
| West Warwick | 31 | 700 | 4.4% |
| Westerly | 11 | 291 | 3.8% |
| Woonsocket | 71 | 1,308 | 5.4% |
| Total | 1,343 | 25,399 | 5.3% |

| Prevalence >=5 µg/dL by Town for 2014 in Rhode Island | | | |
|---|-------------------------------|---------------------|-----------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source : RI CLPPP and LESS Database as of 4/9/2015 | | | |
| TownName | # Children with BLL>=5mc g/dL | # children Screened | Prevalence Rate |
| Barrington | 7 | 429 | 1.6% |
| Bristol | 19 | 483 | 3.9% |
| Burrillville | 10 | 267 | 3.7% |
| Central Falls | 58 | 850 | 6.8% |
| Charlestown | 2 | 91 | 2.2% |
| Coventry | 20 | 643 | 3.1% |
| Cranston | 54 | 1,691 | 3.2% |
| Cumberland | 11 | 774 | 1.4% |
| East Greenwich | 5 | 308 | 1.6% |
| East Providence | 76 | 1,365 | 5.6% |
| Exeter | 2 | 85 | 2.4% |
| Foster | 3 | 55 | 5.5% |
| Glocester | 4 | 134 | 3.0% |
| Hopkinton | 2 | 127 | 1.6% |
| Jamestown | 2 | 72 | 2.8% |
| Johnston | 13 | 569 | 2.3% |
| Lincoln | 9 | 367 | 2.5% |
| Little Compton | 3 | 44 | 6.8% |
| Middletown | 7 | 347 | 2.0% |
| Narragansett | 4 | 123 | 3.3% |
| New Shoreham | 5 | 23 | 13.0% |
| Newport | 35 | 523 | 6.9% |
| North Kingstown | 3 | 445 | 0.7% |
| North Providence | 21 | 571 | 3.7% |
| North Smithfield | 1 | 196 | 0.5% |
| Pawtucket | 164 | 2,390 | 6.9% |
| Portsmouth | 11 | 278 | 4.0% |
| Providence | 579 | 7,106 | 8.1% |
| Richmond | 6 | 73 | 8.2% |
| Saunderstown | 5 | 146 | 3.4% |
| Smithfield | 4 | 270 | 1.5% |
| South Kingstown | 21 | 413 | 5.1% |
| Tiverton | 10 | 351 | 2.8% |
| Warren | 8 | 272 | 2.9% |
| Warwick | 47 | 1,605 | 2.9% |
| West Greenwich | 1 | 67 | 1.5% |
| West Warwick | 26 | 744 | 3.5% |
| Westerly | 17 | 297 | 5.7% |
| Woonsocket | 64 | 1,263 | 5.1% |
| Total | 1,338 | 25,857 | 5.2% |

Data Source: RI CLPPP LESS database as of 3/10/2015

| Prevalence >= 5 µg/dL by Town for 2013 in Rhode Island | | | |
|---|------------------------------|---------------------|--------------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source: RI CLPPP and LESS Database as of 5/30/2014 | | | |
| TownName | Confirmed >= 5 in 2013 | Screened in 2013 | Prevalence in 2013 |
| Barrington | 10 | 455 | 2.2% |
| Bristol | 17 | 499 | 3.4% |
| Burnsville | 15 | 283 | 5.3% |
| Central Falls | 85 | 924 | 9.2% |
| Charlestown | 2 | 105 | 1.9% |
| Coventry | 19 | 567 | 3.4% |
| Cranston | 71 | 1,681 | 4.2% |
| Cumberland | 15 | 714 | 2.1% |
| East Greenwich | 5 | 269 | 2.2% |
| East Providence | 83 | 1,427 | 5.8% |
| Exeter | 5 | 105 | 4.8% |
| Foster | 1 | 69 | 1.4% |
| Glocester | 2 | 127 | 1.6% |
| Hopkinton | 9 | 125 | 7.2% |
| Jamestown | 5 | 60 | 8.3% |
| Johnston | 12 | 557 | 2.2% |
| Lincoln | 12 | 387 | 3.1% |
| Little Compton | 5 | 54 | 5.6% |
| Middletown | 13 | 384 | 3.4% |
| Narragansett | 2 | 129 | 1.6% |
| New Shoreham | 0 | 22 | 0.0% |
| Newport | 40 | 577 | 6.9% |
| North Kingstown | 10 | 492 | 2.0% |
| North Providence | 12 | 619 | 1.9% |
| North Smithfield | 0 | 177 | 0.0% |
| Pawtucket | 169 | 2,434 | 6.9% |
| Portsmouth | 8 | 291 | 2.7% |
| Providence | 648 | 7,347 | 8.8% |
| Richmond | 3 | 108 | 2.8% |
| Slatute | 6 | 185 | 3.2% |
| Smithfield | 1 | 288 | 0.3% |
| South Kingstown | 22 | 462 | 4.8% |
| Tiverton | 9 | 374 | 2.4% |
| Unknown | 0 | 2 | 0.0% |
| Warren | 24 | 267 | 8.4% |
| Warwick | 44 | 1,521 | 2.9% |
| West Greenwich | 3 | 102 | 2.9% |
| West Warwick | 30 | 754 | 4.0% |
| Westerly | 9 | 342 | 2.6% |
| Woonsocket | 35 | 1,226 | 2.9% |
| Total | 1,460 | 26,531 | 5.5% |

| Prevalence >=5 µg/dL by Town for 2012 in Rhode Island | | | |
|--|-----------------------------|---------------------|--------------------|
| For Children <= 72 months of Age at the time of test | | | |
| Data Source : RI CLPPP and LESS Database as of 4/29/2013 | | | |
| TownName | Confirmed >=5 in 2012 | Screened in 2012 | Prevalence in 2012 |
| Barrington | 4 | 441 | 0.9% |
| Bristol | 15 | 524 | 2.9% |
| Burrillville | 16 | 291 | 5.5% |
| Central Falls | 115 | 914 | 12.6% |
| Charlestown | 1 | 119 | 0.8% |
| Coventry | 12 | 568 | 2.1% |
| Cranston | 62 | 1,656 | 3.7% |
| Cumberland | 9 | 786 | 1.1% |
| East Greenwich | 4 | 275 | 1.5% |
| East Providence | 26 | 1,521 | 1.7% |
| Exeter | 0 | 85 | 0.0% |
| Foster | 1 | 59 | 1.7% |
| Glocester | 5 | 142 | 3.5% |
| Hopkinton | 6 | 140 | 4.3% |
| Jamestown | 0 | 63 | 0.0% |
| Johnston | 13 | 564 | 2.3% |
| Lincoln | 16 | 396 | 2.0% |
| Little Compton | 1 | 59 | 1.7% |
| Middletown | 2 | 409 | 0.5% |
| Narragansett | 0 | 124 | 0.0% |
| New Shoreham | 1 | 25 | 4.0% |
| Newport | 29 | 606 | 4.8% |
| North Kingstown | 5 | 475 | 1.1% |
| North Providence | 10 | 567 | 1.8% |
| North Smithfield | 4 | 191 | 2.1% |
| Pawtucket | 151 | 2,526 | 6.0% |
| Portsmouth | 1 | 358 | 0.3% |
| Providence | 1,012 | 7,433 | 13.6% |
| Richmond | 5 | 87 | 5.7% |
| Saunderstown | 2 | 173 | 1.2% |
| Smithfield | 5 | 282 | 1.8% |
| South Kingstown | 9 | 426 | 2.1% |
| Tiverton | 4 | 375 | 1.1% |
| Unknown | 0 | 1 | 0.0% |
| Warren | 8 | 307 | 2.6% |
| Warwick | 35 | 1,534 | 2.3% |
| West Greenwich | 0 | 106 | 0.0% |
| West Warwick | 33 | 768 | 4.3% |
| Westerly | 21 | 341 | 6.2% |
| Woonsocket | 85 | 1,348 | 6.3% |
| Total | 1,720 | 27,065 | 6.4% |